

Evaluation of Bio-Efficiency and Phytotoxicity of Pendimethalin 38.7% Cs (Stomp) in Cotton

Rajamani Raman*, M. Ramachandran and R. Krishnamoorthy

Faculty of Agriculture, Department of Agronomy, Annamalai University, India

*Corresponding author: Raman R, Faculty of Agriculture, Department of Agronomy, Annamalai University, Annamalai Nagar, Chidambaram - 608002, Tamil Nadu, India, Tel: 04144 238 282; Email: ramanagron@gmail.com

Received: November 19, 2021; Accepted: December 21, 2021; Published: December 28, 2021

Abstract

A field experiments were conducted at the Experimental Farm, Department of Agronomy, Annamalai University, Annamalai Nagar to study the evaluation of Pendimethalin 38.7% CS and various cotton herbicides in cotton during August 2018 to January 2019 and August 2009 to January 2010 with cv. MCU 7. The experiment consists of eight treatments viz., Pendimethalin 38.7% CS @ 438.75 g a.i/ha, Pendimethalin 38.7% CS @ 580.50 g a.i/ha, Pendimethalin 38.7% CS @ 677.25 g a.i/ha, Fluchloralin 45% EC @ 1125 g a.i/ha, Pendimethalin 30% EC @ 750 g a.i/ha, Trifluralin 48% EC @ 960 g a.i/ha, Fluchloralin 45% EC @ 2000 g a.i/ha and unweeded check. Regardless of season, all the treatments were found to significantly influenced the weed biometrics, growth, yield components and yield of cotton. Among the treatments tested, Pendimethalin 38.7 % CS @ 677.25 g a.i/ha recorded least weed counts, weed dry matter production, highest weed control index and weed control efficiency favouring higher growth, yield attributes and seed cotton yield in first and second seasons, respectively. This was followed by Pendimethalin 38.7% CS @ 580.50 g a.i/ha. These treatments were significantly superior to the rest in reducing the weed infestation and ultimately increasing seed cotton yield in both seasons. Unweeded check recorded the highest weed counts, weed biomass resulting in the least growth and seed cotton yield in first and second seasons, respectively. From the study, it may be concluded that application of Pendimethalin 38.7% CS @ 677.25 g a.i/ha is considered to be judicious recommendation to cotton farmers in view of inadequate labor and higher manual weeding cost.

Keywords: Cotton; Herbicides; Phytotoxicity; Yield

1. Introduction

Cotton the “white gold” is the one among the most important commercial fiber crop as well as the cash crop of India. It contributes about 85% of raw materials for manufacture of textiles in the world [1]. Globally it is cultivated over an area of 34.6 million hectare with the production of 25.1 million tonnes in world 2008-2009 (Cotton Corporation of India, 2009). India

is the second largest cotton growing country in the world next to China, produces around 16% of the world's total production. The area under cotton is 93.73 lakh hectares, production of 290 lakh bales with the productivity of 526 kg/ha. In Tamil Nadu, it is grown in an area of 1.20 lakh hectares with a production of 5 lakh bales and the productivity of 708 kg/ha. Weeds cause enormous losses and suffering to human beings by way of reduction in crop yield and quality, wastage of human energy and resources and increased cost of cultivation [2]. Pendimethalin, Trifluralin and Fluchloralin is also gaining momentum among cotton growers. Among these herbicides, Pendimethalin 33% is a direct dinitroaniline herbicides commonly used to control grasses and small seeded dicot weed species as pre-emergence (PRE) or pre-plant incorporation (PPI). Gill et al., [3] noted that when Pendimethalin applied on dry land and irrigated crop immediately, exhibited 81% weed control and increased seed cotton yield as compared to delayed irrigation. This study was undertaken to evaluate Pendimethalin 38.7% CS to control weeds in cotton and to find out its phytotoxic effect in cotton crop.

2. Materials and Methods

The experiment was conducted at the Agriculture Experimental farm, Department of Agronomy, Annamalai University, Tamil Nadu during August 2018 to January 2019 and August 2009 to January 2010. The Experimental farm is situated at 11°24'N latitude, 79°44'E longitude and at an altitude of +5.79 meters above mean sea level. The soil of the experimental site was clay loam in texture with 258.5 kg/ha available N, 17.3 kg/ha P and 310.0 kg/ha K and 7.4 pH and 1.20 dS/m EC at the start of the experiment. The experiment was laid out in randomized block design with 3 replications. There were 7 weed control treatments, viz. Pendimethalin 38.7% CS @ 438.75 g a.i/ha, Pendimethalin 38.7% CS @ 580.50 g a.i/ha, Pendimethalin 38.7% CS @ 677.25 g a.i/ha, Fluchloralin 45% EC @ 1125 g a.i/ha, Pendimethalin 30% EC @ 750 g a.i/ha, Trifluralin 48% EC @ 960 g a.i/ha, Fluchloralin 45% EC @ 2000 g a.i/ha and unweeded check. The seeds of cotton were dibbled with a spacing of 60 cm between rows and 30 cm between plants at a depth of 2 cm to 3 cm and then covered with the soil. The herbicides Pendimethalin, Fluchloralin and Trifluralin applied @ 1.5 lit a.i/ha were mixed with water @ 500 lit required doses. These pre-emergence herbicides were sprayed three days after sowing of the cotton seeds by a knapsack sprayer uniformly over the soil. The crop was raised by using recommended package of practices. The seed cotton was harvested from the well opened bolls from 110 DAS onwards and seven pickings were given for the first crop, whereas only four pickings were given for second crop at weekly intervals. The harvested seed cotton was shade dried on a layer of river sand and weighed at uniform moisture content. Economics of cotton was calculated by using prevailing market prices of inputs and outputs under different weed-control treatments.

3. Results and Discussion

3.1 Weed count

The major weed flora observed in cotton field consisted of *Cyperus rotundus*, *Trianthema portulacastrum*, *Phyllanthus maderaspatensis*, *Corchorus trilocularis*, *Convolvulus arvensis*, *Eclipta alba*, *Cleome viscosa*, *Tridax procumbens*, *Echinochloa colonum* and *Cynodan dactylon*. All the treatments were effective in controlling the weeds and caused significant reduction in their population. Among the various herbicides, Pendimethalin 38.7% CS @ 677.25 g a.i/ha was very effective and significantly superior to the rest of the treatments in respect of weed count in both seasons. Effective weed control measures such as pre-emergence application of Pendimethalin at 3 DAS might have helped in minimizing weed population and by reducing weed competition for light, moisture and nutrients as compared to all other herbicide treatments. Weed density were reduced further due to increase in the rates of Pendimethalin application. Pendimethalin, irrespective of rates of application,

provided better weed control than other herbicides. Pendimethalin 30 EC also reduced the weed count, and this result was in consonance with the findings of Naqib Ullabkhan et al., [4]. The highest weed population was recorded under the unweeded check treatment (TABLE 1).

TABLE 1. Effect of Weed-Control Treatments on Weed Count (Nos./m²) in Cotton.

Treatment	30 DAS		60 DAS		90 DAS	
	First year	Second year	First year	Second year	First year	Second year
Pendimethalin 38.7% CS @ 438.75 g a.i/ha	20.57 (4.59)	26.22 (5.17)	24.53 (5.01)	31.64 (5.66)	29.80 (5.17)	42.42 (6.53)
Pendimethalin 38.7% CS @ 580.50 g a.i/ha	13.25 (3.70)	18.81 (4.39)	16.71 (4.14)	22.88 (4.77)	21.07 (4.64)	31.08 (5.61)
Pendimethalin 38.7% CS @ 677.25 g a.i/ha	10.13 (3.26)	15.65 (4.02)	13.49 (3.74)	18.69 (4.38)	17.09 (4.20)	25.91 (5.13)
Fluchloralin 45% EC @ 1125 g a.i/ha	27.85 (5.32)	33.22 (5.80)	31.65 (5.67)	41.67(6.49)	40.44 (6.39)	56.37 (7.54)
Pendimethalin 30% EC @ 750 g a.i/ha	17.35 (4.22)	22.99 (4.84)	21.02 (4.63)	26.92 (5.24)	25.70 (5.11)	36.22 (6.05)
Trifluralin 48% EC @ 960 g a.i/ha	31.95 (5.10)	37.31 (6.14)	35.98 (6.03)	46.69 (6.86)	47.54 (6.93)	63.42 (7.99)
Fluchloralin 45% EC @ 2000 g a.i/ha	24.13 (4.96)	29.67 (5.49)	28.15 (5.35)	36.52 (6.08)	35.03 (5.93)	49.55 (7.08)
Unweeded check	35.83 (6.02)	40.97 (6.43)	40.15 (6.37)	51.58 (7.21)	55.97 (7.49)	71.19 (8.46)
Sem ±	0.10	0.12	0.22	0.25	0.35	0.37
CD (P=0.05)	0.21	0.25	0.45	0.51	0.71	0.75

3.2 Weed dry matter

The weed control treatments significantly influenced the total weed dry matter during both the years. Pendimethalin 38.7% CS @ 677.25 g a.i/ha recorded the lowest weed DMP in both the seasons. This treatment was superior to the other herbicides tested

as it controlled grasses, sedges, and broad-leaved weeds in the critical stages. This might be due to restriction of emergence of fresh weeds in the later stages. Effective control of weeds, due to blocking of their perennation process and inhibition of their germination and growth might have resulted in lesser weed DMP [5]. Lesser weed DMP in this treatment were due to better control of weeds especially in the early stages of crop growth. Unweeded check recorded the highest weeds DMP than other weed control treatment. These findings are in a great agreement with the work of Panwar *et al.*, [6] who reported the efficacy of Pendimethalin for controlling weeds in cotton. The findings of Hiremath and Rao [7] further support the present findings. Richardson *et al.*, [8] who reported that the weeds are controlled more efficiently by the application of Pendimethalin (TABLE 2).

TABLE 2. Effect of weed-control treatments on weed dry matter production (kg/ha) in cotton.

Treatment	30 DAS		60 DAS		90 DAS	
	First year	Second year	First year	Second year	First year	Second year
Pendimethalin 38.7% CS @ 438.75 g a.i/ha	140.30	155.93	179.88	209.22	261.96	333.23
Pendimethalin 38.7% CS @ 580.50 g a.i/ha	90.56	101.19	118.42	137.44	166.54	200.90
Pendimethalin 38.7% CS @ 677.25 g a.i/ha	56.48	61.03	73.12	86.03	92.09	115.27
Fluchloralin 45% EC @ 1125 g a.i/ha	187.20	212.57	244.65	284.86	359.59	494.88
Pendimethalin 30% EC @ 750 g a.i/ha	125.18	140.31	163.67	189.06	241.65	287.58
Trifluralin 48% EC @ 960 g a.i/ha	217.36	253.52	290.63	338.54	434.61	515.55
Fluchloralin 45% EC @ 2000 g a.i/ha	172.95	196.18	225.53	261.25	332.61	414.01
Unweeded check	285.49	325.68	357.50	419.55	534.87	604.52
Sem ±	16.03	19.01	22.60	25.70	37.55	44.30
CD (P=0.05)	32.06	38.03	45.27	51.41	75.11	88.63

3.3 Seed cotton yield

All the treatments had a significant impact on the yield attributes of cotton. Among the herbicides tested, Pendimethalin 38.7 % CS @ 677.25 g a.i/ha registered maximum yield and yield components than other herbicide treatments. It could be attributed to significantly lower weed population, dry matter accumulation of weeds and hence higher number of sympodial branches, number of monopodial branches and number of bolls per plant in this treatment. The superior performance of Pendimethalin might be attributed to reduce crop-weed competition in the initial stages which helped in synchronization of number of sympodial branches, number of monopodial branches and number of bolls per plant. The lowest boll weight was recorded in weedy check by weak plants suppressed by high weed infestation. These are in line with the findings of Gill *et al.*, [3], Hassan *et al.*, [9] and Khan *et al.*, [10]. The increase in yield might be due to effective control of weeds which reduced the crop-weed competition by effective weed control measures such as pre-emergence application of Pendimethalin at 3 DAS offering efficient and prolonged weed control might have favoured the crop with better rooting, higher LAI, pre and post flowering photosynthesis and yield attributes. Severe competition of weeds in unweeded check results in suppression of crop growth and there by reduction in seed cotton yield in both seasons (TABLE 3).

TABLE 3. Effect of weed-control treatments on yield and yield attributes of cotton.

Treatment	No. of monopodial branches/plant		No. of sympodial branches/plant		No. of bolls/plant		Boll weight (g)		Seed cotton yield (kg/ha)	
	First year	Second year	First year	Second year	First year	Second year	First year	Second year	First year	Second year
Pendimethalin 38.7% CS @ 438.75 g a.i/ha	4.12	3.59	19.73	18.29	26.86	25.57	3.59	3.53	1778	1706
Pendimethalin 38.7% CS @ 580.50 g a.i/ha	5.0	4.29	22.17	21.36	30.85	29.47	3.84	3.69	2032	1933
Pendimethalin 38.7% CS @ 677.25 g a.i/ha	5.42	4.62	23.26	22.46	32.84	31.29	3.97	3.78	2160	2035
Fluchloralin 45% EC @ 1125 g a.i/ha	3.25	2.92	16.68	15.49	23.21	21.85	3.33	3.25	1538	1473
Pendimethalin 30% EC @ 750 g a.i/ha	4.58	3.95	21.05	19.21	28.75	27.62	3.68	3.63	1897	1821
Trifluralin 48% EC @ 960 g a.i/ha	2.83	2.56	15.33	14.34	21.76	20.40	3.24	3.16	1433	1356

Fluchloralin 45% EC @ 2000 g a.i/ha	3.68	3.25	18.27	16.94	24.89	23.95	3.43	3.33	1660	1590
Unweeded check	2.38	2.21	14.13	13.13	19.66	18.75	3.09	3.03	1318	1236
Sem ±	0.19	0.15	0.49	0.42	0.57	0.54	0.035	0.025	50.39	49
CD (P=0.05)	0.41	0.32	0.99	0.87	1.14	1.08	0.07	0.05	101	98

3.4 Economics

Among the weed control treatments, the highest gross income was registered with Pendimethalin 38.7% CS @ 677.25 g a.i/ha in both the seasons compared to other treatments. Increased seed cotton yield due to application of Pendimethalin, efficient weed control measures in right time with the right method might have been responsible for increased return per rupee invested in the best treatment (TABLE 4). All weed control treatments had higher gross and net returns than the weedy check during both the years. The benefit: cost ratio under different treatments varied from 1.95 - 2.96 as compared to 1.83 under weedy check. It may be concluded from this study that weeds from cotton can be controlled effectively by using Pendimethalin 38.7% CS @ 677.25 g a.i/ha.

TABLE 4. Effect of Weed-Control Treatments on Economics of Cotton.

Treatment	Cost of cultivation (Rs/ha)		Gross income (Rs/ha)		Net income (Rs/ha)		Benefit: cost ratio	
	First year	Second year	First year	Second year	First year	Second year	First year	Second year
Pendimethalin 38.7% CS @ 438.75 g a.i/ha	20307.74	20307.74	49784	47768	29476.26	27460.26	2.45	2.35
Pendimethalin 38.7 % CS @ 580.50 g a.i/ha	20371.53	20371.53	56896	54124	36524.47	33752.47	2.79	2.65
Pendimethalin 38.7% CS @ 677.25 g a.i/ha	20415.07	20415.07	60480	56980	40064.93	36564.93	2.96	2.79
Fluchloralin 45% EC @ 1125 g a.i/ha	20751.56	20751.56	43064	41244	22312.44	20492.44	2.07	1.98
Pendimethalin 30% EC @ 750 g a.i/ha	20455.31	20455.31	53116	50988	32660.69	30532.69	2.59	2.49

Trifluralin 48% EC @ 960 g a.i/ha	20570.39	20570.39	40404	17968	19553.61	17397.61	1.95	1.84
Fluchloralin 45% EC @ 2000 g a.i/ha	21250.31	21250.31	46480	44520	25229.69	23269.69	2.18	2.09
Unweeded check	20110.31	20110.31	36904	34608	16793.69	14497.69	1.83	1.72

REFERENCES

1. Muhammad D, Afzal MN, Raza I, et al. Growth and development of cotton (*Gossypium hirsutum* L.) as affected by different methods of Pendimethalin application. *Pak J Weed Sci Res.* 2009;15(1):11-7.
2. Singh G. Weed management for sustainable agriculture. National Biennial Conference, ISWS, Ludhiana, Punjab, pp. 1-6. 2005.
3. Gill MI, Anwar M, Mohammad D. Effect of pre-planting irrigation (rouni) and time of pre-emergence application of Stomp 330 E (Pendimethalin) in cotton. Paper read at the 5th Pakistan Weed Science Conference March 1996, NARC, Islamabad. 1996.
4. Khan NU, Khan SU, Hassan G, et al. Studies on weed control in cotton (*Gossypium hirsutum* L.). *Online J Biol Sci.* 2001;1(3):143-5.
5. Hadizadeh MHS, Noroozadeh S, Rahimion H. Effect of row spacing and weed free period on yield components of cotton (*Gossypium hirsutum* L.). *Applied Entomol Phytopathal.* 2002;69(2):171-87.
6. Panwar RS, Malik RS, Rathi SS, et al. Chemical weed control in cotton. *Ind J Weed Sci.* 2001;33(1&2):14-7.
7. Hirematha KA, Sathyanarayann Rao. Chemical weed control studies in irrigated hybrid cotton (*Gossypium hirsutum* L.). *Crop Res.* 2001;21(1):41-5.
8. Richardson RJ, Wilson HP, Hines TE. Pre-emergence herbicides followed by trifloxysulfuron post-emergence in cotton. *Weed Tech.* 2007;21(1):1-6.
9. Hassan G, Khan NU, Nawaz Q. Chemical control of weeds in cotton (*G. hirsutum* L.) under D.I. Khan Conditions. Paper read at the 1st Crop Protection Conference (CAP) April 21-23, 1996, NWFP Agricultural University Peshawar, Pakistan, Abstracts.pp.41. 1996.
10. Khan NU, Khan SU, Hassan G, et al. Studies on weed control in cotton (*G. hirsutum* L.). *Online J Biol Sci.* 2001;1(3):143-5.